

ON THE CYCLE STRUCTURE OF REPEATED EXPONENTIATION MODULO A PRIME POWER

MIN SHA

ABSTRACT. We obtain some results about the repeated exponentiation modulo a prime power from the viewpoint of arithmetic dynamical systems. Especially, we extend two asymptotic formulas about periodic points and tails in the case of modulo a prime to the case of modulo a prime power.

1. INTRODUCTION

For a positive integer M , denote by $\mathbb{Z}/M\mathbb{Z}$ the residue ring of $\mathbb{Z}$ modulo M and $(\mathbb{Z}/M\mathbb{Z})^*$ the unit group. For an integer $k \geq 2$, we consider the following endomorphism of $(\mathbb{Z}/M\mathbb{Z})^*$,

$$f : (\mathbb{Z}/M\mathbb{Z})^* \rightarrow (\mathbb{Z}/M\mathbb{Z})^*, x \rightarrow x^k.$$

For any initial value $x \in (\mathbb{Z}/M\mathbb{Z})^*$, we repeat the action of f , then we get a sequence

$$x_0 = x, x_n = x_{n-1}^k, n = 1, 2, 3, \dots$$

This sequence is known as the power generator of pseudorandom numbers. Studying such sequences in the cases that M is a prime or a product of two distinct primes, is of independent interest and is also important for several cryptographic applications, see [1, 6]. From the viewpoint of cryptography, there are numerous results about these sequences, see the papers mentioned in [2], more recently see [3] and its references.

If we view $(\mathbb{Z}/M\mathbb{Z})^*$ as a vertex set and draw a directed edge from a to b if $f(a) = b$, then we get a digraph. There are also many results in this direction, see [12] and the papers mentioned there, more recently see [8, 9, 10, 11].

As [2], in this article we will study $(\mathbb{Z}/M\mathbb{Z})^*$ under the action of f from the viewpoint of arithmetic dynamical systems, where M is a prime power. Especially we will extend two asymptotic formulas in [2] to the case of modulo a prime power.

It is easy to see that for any initial value $x \in (\mathbb{Z}/M\mathbb{Z})^*$ the corresponding sequence becomes eventually periodic, that is, for some positive integer $s_{k,M}(x)$ and *tail* $t_{k,M}(x) < s_{k,M}(x)$, the elements $x_0 = x, x_1, \dots, x_{s_{k,M}(x)-1}$ are pairwise distinct and $x_{s_{k,M}(x)} = x_{t_{k,M}(x)}$. So we can define a *tail function* $t_{k,M}$ on $(\mathbb{Z}/M\mathbb{Z})^*$.

The sequence $x_{t_{k,M}(x)}, \dots, x_{s_{k,M}(x)-1}$, ordered up to a cyclic shift, is called a *cycle*. The *cycle length* is $c_{k,M}(x) = s_{k,M}(x) - t_{k,M}(x)$. The elements in the cycle are called *periodic points* and their *periods* are $c_{k,M}(x)$. So we can define a *cycle length function* $c_{k,M}$ on $(\mathbb{Z}/M\mathbb{Z})^*$. In particular, [4, 5] gave lower bounds for the largest period.

2010 *Mathematics Subject Classification.* Primary 37P35; Secondary 11K45, 11B50.

Key words and phrases. Repeated powering, Periodic point, Tail, Primes in arithmetic progressions.

We denote by $P_r(k, M)$ and $P(k, M)$ respectively the number of periodic points with period r and the number of periodic points in $(\mathbb{Z}/M\mathbb{Z})^*$. Also, we denote by $C_r(k, M)$ and $C(k, M)$ respectively the number of cycles with length r and the number of cycles in $(\mathbb{Z}/M\mathbb{Z})^*$. We denote the average values of $c_{k,M}(x)$ and $t_{k,M}(x)$ over all $x \in (\mathbb{Z}/M\mathbb{Z})^*$ by $c(k, M)$ and $t(k, M)$ respectively,

$$c(k, M) = \frac{1}{\varphi(M)} \sum_{x \in (\mathbb{Z}/M\mathbb{Z})^*} c_{k,M}(x), \quad t(k, M) = \frac{1}{\varphi(M)} \sum_{x \in (\mathbb{Z}/M\mathbb{Z})^*} t_{k,M}(x),$$

where φ is the Euler totient function.

When M is an odd prime power, we will derive explicit formulas for $P_r(k, M)$ and $C_r(k, M)$ by the results in [10], and we will also derive explicit formulas for $c(k, M)$ and $t(k, M)$ which generalize those in [11].

For two integers $r, m \geq 1$, we call the limit of $\lim_{X \rightarrow \infty} \frac{1}{\pi(X)} \sum_{p \leq X} P_r(k, p^m)$ the asymptotic mean number of periodic points with period r in $(\mathbb{Z}/p^m\mathbb{Z})^*$ for different choices of prime p , and we denote it by $AP_r(k, m)$. Similarly, we can define the asymptotic mean number for cycles with length r and denote it by $AC_r(k, m)$. We will derive explicit formulas for $AP_r(k, m)$ and $AC_r(k, m)$.

For an integer $m \geq 1$, following [11], we study the average values of $P(k, p^m)$ and $t(k, p^m)$ over all primes $p \leq N$,

$$S_0(k, m, N) = \frac{1}{\pi(N)} \sum_{p \leq N} P(k, p^m), \quad S(k, m, N) = \frac{1}{\pi(N)} \sum_{p \leq N} t(k, p^m).$$

where, as usual, $\pi(N)$ is the number of primes $p \leq N$. Following the method in [2], we will get asymptotic formulas for $S_0(k, m, N)$ and $S(k, m, N)$.

2. PREPARATIONS

For two integers l and n , we denote their greatest common divisor by $\gcd(l, n)$. For a positive integer n , we denote by $\tau(n)$ the number of its positive divisors. Theorem 4.9 in [7] tells us that

$$(2.1) \quad \lim_{X \rightarrow \infty} \frac{1}{\pi(X)} \sum_{p \leq X} \gcd(p-1, n) = \tau(n).$$

For two integers $m \geq 1$ and $n \geq 2$, we denote the largest prime divisor of n by q . Then we have

$$(2.2) \quad \begin{aligned} & \lim_{X \rightarrow \infty} \frac{1}{\pi(X)} \sum_{p \leq X} \gcd(p^{m-1}(p-1), n) \\ &= \lim_{X \rightarrow \infty} \frac{1}{\pi(X)} \sum_{q < p \leq X} \gcd(p^{m-1}(p-1), n) \\ &= \lim_{X \rightarrow \infty} \frac{1}{\pi(X)} \sum_{q < p \leq X} \gcd(p-1, n) \\ &= \tau(n). \end{aligned}$$

Notice that if p is an odd prime, $\gcd(p^m - p^{m-1}, n)$ is the number of solutions of the equation $x^n = 1$ in $(\mathbb{Z}/p^m\mathbb{Z})^*$.

Given two integers a and n with $\gcd(a, n) = 1$, following the method in the proof of Formula (2) in [2], we can get

$$(2.3) \quad \sum_{\substack{p \leq X \\ p \equiv a \pmod{n}}} p^m = \frac{X^{m+1}}{(m+1)\varphi(n)\ln X} + O(X^{m+1}\ln^{-2}X).$$

Then we have

$$(2.4) \quad \sum_{\substack{p \leq X \\ p \equiv a \pmod{n}}} p^{m-1}(p-1) = \frac{X^{m+1}}{(m+1)\varphi(n)\ln X} + O(X^{m+1}\ln^{-2}X).$$

Following the same method in the proof of Formula (4) in [2], we have

$$(2.5) \quad \sum_{\substack{p \leq X \\ p \equiv a \pmod{n}}} p^{m-1}(p-1) = O\left(\frac{X^{m+1}}{n} + X^m\right).$$

3. MAIN RESULTS

For two integers d and n satisfying $\gcd(d, n) = 1$, we denote the multiplicative order of n modulo d by $\text{ord}_d n$. For an integer n and a prime p , we denote $v_p(n)$ the exact power of p dividing n .

Let μ be the Möbius function. For a real number a , we denote $\lceil a \rceil$ the least integer which is not less than a .

Write $k = p_1^{n_1} p_2^{n_2} \cdots p_s^{n_s} \geq 2$, where $p_1, \dots, p_s$ are distinct primes, $p_1 < p_2 < \cdots < p_s$ and $n_1, \dots, n_s \geq 1$. Let m be a fixed positive integer.

Proposition 3.1. *Let p be an odd prime and r be a positive integer. Write $p^m - p^{m-1} = p_1^{r_1} \cdots p_s^{r_s} \cdot \rho$, where $r_1, \dots, r_s \geq 0$ are integers and $\gcd(p_1 \cdots p_s, \rho) = 1$. We have*

- (1) $C_r(k, p^m) = \frac{1}{r} \sum_{d|r} \mu(d) \gcd(p^m - p^{m-1}, k^{r/d} - 1).$
- (2) $P_r(k, p^m) = \sum_{d|r} \mu(d) \gcd(p^m - p^{m-1}, k^{r/d} - 1).$
- (3) $P(k, p^m) = \rho.$
- (4) $C(k, p^m) = \sum_{d|\rho} \frac{\varphi(d)}{\text{ord}_d k}.$
- (5) For any $x \in (\mathbb{Z}/p^m \mathbb{Z})^*$, denote $\text{ord}_{p^m} x$ by $\text{ord} x$, $c_{k, p^m}(x) = \text{ord}_{\gcd(\text{ord} x, \rho)} k.$
- (6) $c(k, p^m) = \frac{1}{\rho} \sum_{d|\rho} \varphi(d) \text{ord}_d k.$
- (7) For any $x \in (\mathbb{Z}/p^m \mathbb{Z})^*$, denote $\text{ord}_{p^m} x$ by $\text{ord} x$,

$$t_{k, p^m}(x) = \max \left\{ \left\lceil \frac{v_{p_1}(\text{ord} x)}{n_1} \right\rceil, \left\lceil \frac{v_{p_2}(\text{ord} x)}{n_2} \right\rceil, \dots, \left\lceil \frac{v_{p_s}(\text{ord} x)}{n_s} \right\rceil \right\}.$$

$$(8) \quad t(k, p^m) = \frac{1}{p_1^{r_1} \cdots p_s^{r_s}} \sum_{d|p_1^{r_1} \cdots p_s^{r_s}} \varphi(d) \max \left\{ \left\lceil \frac{v_{p_1}(d)}{n_1} \right\rceil, \dots, \left\lceil \frac{v_{p_s}(d)}{n_s} \right\rceil \right\} \dots$$

Proof. (1) and (2) By Möbius inversion formula and Theorem 5.6 in [10].

(3) A special case of Corollary 3 in [12].

(4) By Theorem 2 and Theorem 3 in [12].

(5) By Lemma 3 and Theorem 2 in [12].

(6) Denote $p_1^{r_1} \cdots p_s^{r_s}$ by w , from (5), we have

$$\begin{aligned} c(k, p^m) &= \frac{1}{p^m - p^{m-1}} \sum_{x \in (\mathbb{Z}/p^m \mathbb{Z})^*} c_{k, p^m}(x) \\ &= \frac{1}{p^m - p^{m-1}} \sum_{d|\rho} \sum_{n|w} \varphi(dn) \text{ord}_d k \\ &= \frac{1}{p^m - p^{m-1}} \sum_{n|w} \varphi(n) \sum_{d|\rho} \varphi(d) \text{ord}_d k = \frac{1}{\rho} \sum_{d|\rho} \varphi(d) \text{ord}_d k. \end{aligned}$$

(7) Let w_x be the factor of $\text{ord} x$ such that $\frac{\text{ord} x}{w_x}$ is the largest factor relatively prime to k . By Lemma 3 in [12], we have $t_{k, p^m}(x)$ is the least non-negative integer l such that $w_x | k^l$. In other words, $t_{k, p^m}(x)$ is the least non-negative integer l such that $v_{p_i}(\text{ord} x) \leq l n_i$, for any $1 \leq i \leq s$. Then we get the desired result.

(8) Notice that for any $x \in (\mathbb{Z}/p^m \mathbb{Z})^*$, $\text{ord} x | (p^m - p^{m-1})$, and there are $\varphi(\text{ord} x)$ elements with the order $\text{ord} x$. By (7), we have

$$t(k, p^m) = \frac{1}{p^m - p^{m-1}} \sum_{d|(p^m - p^{m-1})} \varphi(d) \max \left\{ \left\lceil \frac{v_{p_1}(d)}{n_1} \right\rceil, \left\lceil \frac{v_{p_2}(d)}{n_2} \right\rceil, \dots, \left\lceil \frac{v_{p_s}(d)}{n_s} \right\rceil \right\}.$$

Furthermore, we have

$$\begin{aligned} t(k, p^m) &= \frac{1}{p^m - p^{m-1}} \sum_{d|p_1^{r_1} \cdots p_s^{r_s} \rho} \varphi(d) \max \left\{ \left\lceil \frac{v_{p_1}(d)}{n_1} \right\rceil, \dots, \left\lceil \frac{v_{p_s}(d)}{n_s} \right\rceil \right\} \\ &= \frac{1}{p^m - p^{m-1}} \sum_{i_1=0}^{r_1} \cdots \sum_{i_s=0}^{r_s} \sum_{d|\rho} \varphi(p_1^{i_1} \cdots p_s^{i_s} d) \max \left\{ \left\lceil \frac{i_1}{n_1} \right\rceil, \dots, \left\lceil \frac{i_s}{n_s} \right\rceil \right\} \\ &= \frac{1}{p^m - p^{m-1}} \sum_{d|\rho} \varphi(d) \sum_{i_1=0}^{r_1} \cdots \sum_{i_s=0}^{r_s} \varphi(p_1^{i_1} \cdots p_s^{i_s}) \max \left\{ \left\lceil \frac{i_1}{n_1} \right\rceil, \dots, \left\lceil \frac{i_s}{n_s} \right\rceil \right\} \\ &= \frac{1}{p_1^{r_1} \cdots p_s^{r_s}} \sum_{d|p_1^{r_1} \cdots p_s^{r_s}} \varphi(d) \max \left\{ \left\lceil \frac{v_{p_1}(d)}{n_1} \right\rceil, \dots, \left\lceil \frac{v_{p_s}(d)}{n_s} \right\rceil \right\}. \end{aligned}$$

□

Remark 3.2. If we put $k = 2$ and $m = 1$, then the formulas (3),(4),(6) and (8) correspond to Theorem 6 in [11].

Remark 3.3. Since the conclusions in [10] and [12] we apply are about the general case of modulo a positive integer, it is easy to get similar formulas for the case of $p = 2$.

Proposition 3.4. Let r be a positive integer, we have

$$(3.1) \quad AP_r(k, m) = \sum_{d|r} \mu(d) \tau(k^{r/d} - 1),$$

$$(3.2) \quad AC_r(k, m) = \frac{1}{r} \sum_{d|r} \mu(d) \tau(k^{r/d} - 1).$$

Proof. Combing (2.2) and Proposition 3.1 (1) and (2), we can get the desired formulas. □

In the following, we denote by Ω the set of positive $\mathcal{S}$ -units with $\mathcal{S} = \{p_1, \dots, p_s\}$. Here a positive $\mathcal{S}$ -unit means a positive integer whose prime divisors all belong to $\mathcal{S}$.

Proposition 3.5. We have $\lim_{N \rightarrow \infty} \frac{S_0(k, m, N)}{N^m} = \frac{1}{m+1} \left(\prod_{i=1}^s \frac{p_i^2}{p_i^2 - 1} - 1 \right).$

Proof. Put $Q = p_1 p_2 \cdots p_s$ and denote by $\mathcal{U}_Q$ the set of integer $u, 1 \leq u \leq Q$, such that $\gcd(u, Q) = 1$.

For each odd prime p , let ρ_p be the largest divisor of $p^m - p^{m-1}$ coprime to $p_1 p_2 \cdots p_s$. It is easy to see

$$\lim_{N \rightarrow \infty} \frac{S_0(k, m, N)}{N^m} = \lim_{N \rightarrow \infty} \frac{1}{N^m \pi(N)} \sum_{p_s < p \leq N} \rho_p.$$

Notice that if a prime $p > p_s$, then $v_{p_i}(p^m - p^{m-1}) = v_{p_i}(p - 1)$ for any $1 \leq i \leq s$. Hence, following the method in Theorem 2 of [2], we have

$$\lim_{N \rightarrow \infty} \frac{S_0(k, m, N)}{N^m} = \lim_{N \rightarrow \infty} \frac{1}{N^m \pi(N)} \sum_{q \in \Omega} q^{-1} \sum_{u \in \mathcal{U}_Q} \sum_{\substack{p \leq N \\ p \equiv qu + 1 \pmod{qQ}}} (p^m - p^{m-1}).$$

Following the method in Theorem 2 of [2], we have

$$\lim_{N \rightarrow \infty} \frac{S_0(k, m, N)}{N^m} = \frac{1}{m+1} \sum_{q \in \Omega} \frac{1}{q^2}.$$

Moreover, we have

$$\begin{aligned} \sum_{q \in \Omega} \frac{1}{q^2} &= \sum_{i_1, \dots, i_s=0}^{\infty} \frac{1}{(p_1^{i_1} \cdots p_s^{i_s})^2} - 1 \\ &= \sum_{i_1=0}^{\infty} \frac{1}{p_1^{2i_1}} \cdots \sum_{i_s=0}^{\infty} \frac{1}{p_s^{2i_s}} - 1 \\ &= \prod_{i=1}^s \frac{p_i^2}{p_i^2 - 1} - 1. \end{aligned}$$

Hence, we get the desired result. □

Corollary 3.6. *We have $\frac{1}{k^2(m+1)} < \lim_{N \rightarrow \infty} \frac{S_0(k, m, N)}{N^m} < \frac{2^s - 1}{m+1}$.*

Proof. Notice that for any prime p , we have

$$1 + p^{-2} < \frac{p^2}{p^2 - 1} = 1 + \frac{1}{p^2 - 1} < 2.$$

□

Given $q = p_1^{r_1} \cdots p_s^{r_s} \in \Omega$, we denote

$$\psi(q) = \frac{1}{q} \sum_{d|q} \varphi(d) \max \left\{ \left\lceil \frac{v_{p_1}(d)}{n_1} \right\rceil, \dots, \left\lceil \frac{v_{p_s}(d)}{n_s} \right\rceil \right\}.$$

Proposition 3.7. *We have $\lim_{N \rightarrow \infty} S(k, m, N) = \sum_{q \in \Omega} \frac{\psi(q)}{q}$.*

Proof. Given $q = p_1^{r_1} \cdots p_s^{r_s} \in \Omega$. Suppose $r_1 \geq 1$, we want to estimate $\frac{1}{q} \sum_{d|q} \varphi(d) \lceil \frac{v_{p_1}(d)}{n_1} \rceil$.

For simplicity, we replace p_1, r_1 and n_1 by p, r and n respectively. By division algorithm, we write $r = ln + d$ with $0 \leq d < n$. We have

$$\begin{aligned}
\frac{1}{q} \sum_{d|q} \varphi(d) \lceil \frac{v_p(d)}{n} \rceil &= \frac{1}{p^r} \sum_{d|p^r} \varphi(d) \lceil \frac{v_p(d)}{n} \rceil \\
&= \frac{p-1}{p^r} \sum_{i=1}^r p^{i-1} \lceil \frac{i}{n} \rceil \\
&= \frac{p-1}{p^r} \left[\sum_{i=1}^n p^{i-1} + \sum_{i=n+1}^{2n} 2p^{i-1} + \cdots + \sum_{i=(l-1)n+1}^{ln} lp^{i-1} + \sum_{i=ln+1}^{ln+d} (l+1)p^{i-1} \right] \\
&= \frac{p^n-1}{p^r} [1 + 2p^n + \cdots + lp^{(l-1)n}] + \frac{(l+1)p^{ln}(p^d-1)}{p^r} \\
&= \frac{lp^{ln}}{p^r} - \frac{p^{ln}-1}{p^r(p^n-1)} + \frac{(l+1)p^{ln}(p^d-1)}{p^r} \\
&\leq l + (l+1) \leq 3r.
\end{aligned}$$

Hence, we have

$$\begin{aligned}
\psi(q) &\leq \frac{1}{q} \sum_{d|q} \varphi(d) \left(\lceil \frac{v_{p_1}(d)}{n_1} \rceil + \cdots + \lceil \frac{v_{p_s}(d)}{n_s} \rceil \right) \\
(3.3) \quad &\leq 3(r_1 + \cdots + r_s) \\
&\leq \frac{3}{\ln 2} \ln q = O(\ln q).
\end{aligned}$$

Similarly to Proposition 3.5, by Proposition 3.1 (8), we have

$$\lim_{N \rightarrow \infty} S(k, m, N) = \lim_{N \rightarrow \infty} \frac{1}{\pi(N)} \sum_{q \in \Omega} \psi(q) \sum_{u \in \mathcal{U}_Q} \sum_{\substack{p \leq N \\ p \equiv qu+1 \pmod{qQ}}} 1.$$

Then following the method in Theorem 2 of [2], we can get the desired result. $\square$

Corollary 3.8. *We have $\frac{1}{k} < \lim_{N \rightarrow \infty} S(k, m, N) < \frac{5\sqrt{p_1} \cdots \sqrt{p_s}}{(\sqrt{p_1}-1) \cdots (\sqrt{p_s}-1)}$.*

Proof. On one hand we have

$$\begin{aligned}
\sum_{q \in \Omega} \frac{\psi(q)}{q} &> \sum_{i_1 \geq n_1, \dots, i_s \geq n_s} \frac{\varphi(p_1^{i_1} \cdots p_s^{i_s})}{(p_1^{i_1} \cdots p_s^{i_s})^2} \\
&= \frac{(p_1-1) \cdots (p_s-1)}{p_1 \cdots p_s} \sum_{i_1 \geq n_1}^{\infty} \frac{1}{p_1^{i_1}} \cdots \sum_{i_s \geq n_s}^{\infty} \frac{1}{p_s^{i_s}} \\
&= \frac{1}{k}.
\end{aligned}$$

On the other hand, by (3.3) we have $\psi(q) < 5 \ln q$, then we have

$$\begin{aligned}
\sum_{q \in \Omega} \frac{\psi(q)}{q} &< \sum_{q \in \Omega} \frac{5 \ln q}{q} \\
&< 5 \sum_{q \in \Omega} \frac{1}{\sqrt{q}} \\
&= 5 \sum_{i_1=0, \dots, i_s=0}^{\infty} \frac{1}{\sqrt{p_1^{i_1} \cdots p_s^{i_s}}} \\
&= \frac{5\sqrt{p_1} \cdots \sqrt{p_s}}{(\sqrt{p_1}-1) \cdots (\sqrt{p_s}-1)}.
\end{aligned}$$

$\square$

4. REMARKS ON THE GENERAL CASE

In this section, we will give some remarks on the case of modulo a positive integer.

We can deduce formulas for $C_r(k, M)$ and $P_r(k, M)$ directly from Theorem 5.6 in [10]. Corollary 3 in [12] has given a formula for $P(k, M)$. We can also derive a formula for $C(K, M)$ directly by applying Theorem 2 and Theorem 3 in [12].

Following the same methods, we can easily determine the cycle length function $c_{k,M}(x)$ and the tail function $t_{k,M}(x)$ on $(\mathbb{Z}/M\mathbb{Z})^*$, then we can get formulas for $c(k, M)$ and $t(k, M)$.

In fact, [12] and [10] can tell us more information about the properties of repeated exponentiation modulo a positive integer.

5. ACKNOWLEDGMENT

We would like to thank Prof. I.E. Shparlinski for suggesting this problem and for his helpful advice.

REFERENCES

1. L. Blum, M. Blum and M. Shub, *A simple unpredictable pseudo-random number generator*, SIAM J. Comp., **15** (1986), 364-383.
2. W.-S. Chou and I.E. Shparlinski, *On the Cycle Structure of Repeated Exponentiation Modulo a Prime*, J. Number Theory, **107** (2004), 345-356.
3. E. D. El-Mahassni, *On the Distribution of the Power Generator over a Residue Ring for Parts of the Period*, Rev. Mat. Complut., **21** (2008), 319-325.
4. J. B. Friedlander, C. Pomerance and I. E. Shparlinski, *Period of the power generator and small values of Carmichael's function*, Math. Comp., **70** (2001), 1591-1605.
5. P. Kurlberg and C. Pomerance, *On the period of the linear congruential and power generators*, Acta Arith., **119** (2005), 149-169.
6. J. C. Lagarias, *Pseudorandom number generators in cryptography and number theory*, Proc. Symp. in Appl. Math., Amer. Math. Soc., Providence, RI, **42** (1990), 115-143.
7. M. Nilsson, *Cycles of monomials and perturbed monomial p -adic dynamical systems*, Ann. Math. Blaise Pascal, **7**(1) (2000), 37-63.
8. L. Somer, M. Křížek, *Structure of digraphs associated with quadratic congruences with composite moduli*, Discrete Math. **306** (2006) 2174-2185.
9. L. Somer, M. Křížek, *On semiregular digraphs of the congruence $x^k \equiv y \pmod{n}$* , Comment. Math. Univ. Carolin. **48** (2007), 41-58.
10. L. Somer, M. Křížek, *On symmetric digraphs of the congruence $x^k \equiv y \pmod{n}$* , Discrete Math. **309** (2009) 1999-2009.
11. T. Vasiga, J. Shallit, *On the iteration of certain quadratic maps over $GF(p)$* , Discrete Math. **277** (2004) 219-240.
12. B. Wilson, *Power digraphs modulo n* , Fibonacci Quart. **36** (1998) 229-239.

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UNIVERSITÉ BORDEAUX 1, 33405 TALENCE CEDEX, FRANCE

E-mail address: shamin2010@gmail.com